

### Sensivität/Spezifität der kommerziellen Zweistufen-Diagnostik auf Lyme-Borreliose

Studie/Jahr	Sensivität	Spezifität
Schmitz et al, 1993 [1]	66%	100%
Engstrom et al, 1995 [2]	55%	96%
Ledue et al, 1996 [3]	50%	100%
Bakken et al, 1997 [4]	75%	81%
Trevejo et al, 1999 [5]	29%	100%
Nowakowski et al, 2001 [6]	66%	99%
Bacon et al, 2003 [7]	68%	99%
Coulter et al, 2005 [8]	18%	-
Wormser et al, 2008 [9]	14,1%	-
<b>Mittelwert</b>	<b>49,01%</b>	<b>96%</b>

#### Referenzen:

- [1] Schmitz et al. Eur J Clin Microbiol Infect Dis. 1993; 12:419-24
- [2] Engstrom et al. J Clin Microbiol. 1995;33:419-27
- [3] Ledue et al. J Clin Microbiol. 1996;34:2343-50
- [4] Bakken et al. J Clin Microbiol 1997; 35(3):537-543
- [5] Trevejo et al. J Infect Dis. 1999;179:931-8
- [6] Bacon et al. J Infect Dis. 2003;187:1187-99
- [7] Coulter et al. J Clin Microbiol 2005;43:5080-5084
- [8] Wormser et al. Clin Vaccine Immunol. 2008;(10):1519-22